

# COLLECTIVE REVIEWS

## TOURNIQUETS: A REVIEW OF CURRENT USE WITH PROPOSALS FOR EXPANDED PREHOSPITAL USE

Gerard S. Doyle, MD, MPH, Peter P. Taillac, MD

### QUERY SHEET

- Q1: Au: Need received/accepted dates.
- Q2: Au: Pls. provide keywords.
- Q3: Au: Right running head ok?
- Q4: Au: Full page range?
- Q5: Au: City and state of mtg?
- Q6: Au: Please provide URL.

# COLLECTIVE REVIEWS

## TOURNIQUETS: A REVIEW OF CURRENT USE WITH PROPOSALS FOR EXPANDED PREHOSPITAL USE

Gerard S. Doyle, MD, MPH, Peter P. Taillac, MD

5 ABSTRACT

The use of arterial tourniquets in prehospital emergency care has been fraught with controversy and superstition for many years despite the potential utility of these tools. This review examines this controversy in the context of the history of the tourniquet as well as its recent use in surgery and modern battlefield casualty care. Safe prehospital tourniquet use is widespread in the military and is based on sound physiologic data and clinical experience from the surgical use of tourniquets. The physiologic, pathophysiologic, and clinical underpinnings of safe tourniquet use are reviewed here, along with a discussion of alternatives to tourniquets. Prehospital settings in which tourniquets are useful include tactical emergency medical services (EMS) and other law enforcement environments as well as disaster and mass casualty incidents. Beyond this, we present arguments for tourniquet use in more routine EMS settings, in which it may be beneficial but has heretofore been considered

inappropriate. Protocols that foster safe, effective prehospital tourniquet use in these settings are then presented. Finally, we discuss future directions in which tourniquet research and other initiatives will further enhance the safe, rational use of this potentially life-saving tool. **Key words:**

PREHOSPITAL EMERGENCY CARE 2008;12:1-16

### INTRODUCTION

Arterial tourniquets have a long and checkered history. Since their introduction, probably in ancient Roman times, their basic configuration has changed little to this day. They have been called both life saving and "an instrument of the [D]evil that sometimes saves a life."<sup>1</sup> Tourniquets have undergone a dramatic resurgence in popularity in the past decade, owing primarily to an emphasis on rapid hemostasis on the battlefield during recent wars.

Traditionally, tourniquet use has been ruled by the dictum *primum non nocere* or "first, do no harm." Tourniquets have been thought to be dangerous in the hands of prehospital care providers and have usually been seen as a technique of last resort for the emergency medical technician (EMT). This is the result of anecdotal experience from past wars when tourniquets were placed (sometimes unnecessarily) and left in place for extended periods, resulting in limb ischemia, muscle and nerve injury, gangrene, and amputations. However, recent experience with tourniquets in the hands of well-trained military medics, from both the United States and other countries, has resulted in renewed enthusiasm for the instrument in military emergency care. Lives are being saved on modern battlefields as a result of appropriate tourniquet use combined with rapid evacuation of casualties to definitive care. The parallels to modern emergency medical services (EMS) systems are obvious, and it is time to reconsider the tourniquet as a valuable and potentially lifesaving tool for the modern civilian EMT.

Tourniquets, like all medical therapies, have certain dangers inherent in their use. These potential

Q1 Received from the Division of Emergency Medicine, Department of Medicine (GSD) and the Division of Emergency Medicine, Department of Surgery, University of Utah, Salt Lake City, Utah (PPT). Accepted for publication.

Dr Gerard S. Doyle, along with Dr. Peter P. Taillac, developed the concept for this paper. Both authors are members of the University of Utah Division of Emergency Medicine. Each author performed reviews of the relevant literature. Dr. Doyle wrote the first draft of the paper, which was then revised by Dr. Taillac acting independently. Finally, the two edited and revised the paper collaboratively.

The authors wish to acknowledge Dr. David Fosnocht, MD, for his assistance in manuscript review and editing.

None of the contributors had financial support for manuscript development or any financial arrangement that might represent a conflict of interest. Dr. Taillac is Medical Director of the West Valley City, Utah, Fire Department.

As of August 1, 2007, Dr. Doyle has relocated to the University of Wisconsin Division of Emergency Medicine.

Address correspondence and reprint requests to: Gerard S. Doyle, MD, MPH, Division of Emergency Medicine, Department of Medicine, University of Wisconsin Hospital F2/204 CSC, 600 Highland Avenue, Madison, WI 53792. e-mail: gsdoyle@medicine.wisc.edu

doi: 10.1080/10903120801907570

limitations and complications must be addressed prior to increased utilization in civilian EMS. Recent positive military experience with this instrument should not lead to irrational, unfettered tourniquet use. Rather, protocol-driven use by well-trained civilian EMTs would add a valuable weapon to the armamentarium of prehospital emergency providers, as they address difficult problems in controlling extremity hemorrhage.

Recent terrorist activities have served to emphasize the utility of the tourniquet in disaster settings. Disasters, whether man-induced or natural, may result in large numbers of bleeding individuals. Early in such an incident, a few rescue personnel may be required to triage and institute care for a large number of victims. There may not be enough providers to, for example, hold direct pressure on a heavily bleeding wound for long periods. In these circumstances, early utilization of a tourniquet by protocol may be appropriate. Indeed, a primary tourniquet approach, or "tourniquet first," may save lives by allowing the EMT to rapidly stop extremity hemorrhage, move on to other victims, and return later to reassess and possibly remove the tourniquet under circumstances that are more "stable." This is the model that is proving life saving in military field use in Iraq and Afghanistan, and should be considered for civilian EMT use as well.

We present a review of the medical literature focusing on the history, physiology, and complications of tourniquets, their current use, and alternatives to tourniquets for control of bleeding. We then discuss the use of tourniquets in civilian EMS care. Next, we show why EMTs should be facile in the safe use of these tools during situations not well addressed by current protocols. Finally, we argue that civilian EMS as well as law enforcement agencies should adopt expanded indications for tourniquet use under specific protocols.

## History

### Development and Early Uses<sup>2</sup>

The tourniquet arose from the need of battlefield surgeons to control bleeding during surgical amputations, with use being dated as far back as ancient Rome. Ambrose Pare (ca. 1510–1590) is credited with being the first to use the word *tourniquet*, as well as being among the first to record a recommendation for operative use of a tourniquet. He also performed the first-known modification on the tourniquet: A screw was placed over the main vessel of an extremity and tightened with a circumferential strap in place. Around the 17th century, William Fabry and Etienne Morel both used a *windlass*, wherein a stick is used to twist and thus further tighten a constricting band. Many modern designs feature a windlass to allow easy adjustment of tension.

Lister and Esmarch used tourniquets starting in the middle 19th century to introduce bloodless surgery.

Technological advances led Cushing to abandon prior tourniquet designs and introduce a pneumatic tourniquet in 1904. This device made tourniquet application and removal easier, and pressure was more evenly applied to the limb than with prior versions. By the middle of the 20th century, use of tourniquets in extremity surgery to allow operation in a "dry field" was considered routine.

### The Tourniquet in First-Aid

For many generations, tourniquet use in first aid has been controversial. Tourniquets have long been placed in first-aid kits, yet many surgeons who use them enthusiastically in operations have agreed with the tenet that there "is no place for the tourniquet as a first-aid measure."<sup>2</sup> Civilians were felt to be unable to use the instrument safely or effectively, and there has been previous military experience of harm done by inappropriate tourniquet use on bleeding limbs.<sup>3</sup> The most recent American Heart Association and American Red Cross First Aid manuals reflect this philosophy. While recognizing the pivotal role of hemorrhage control, the 2005 First Aid Guidelines of the National First Aid Science Advisory Board recommends only direct pressure and compression dressings (using an elastic bandage) to stop bleeding before EMS providers arrive.<sup>4</sup>

### Military Considerations

The tourniquet has a rich historical tradition in military medicine, in contrast to civilian EMS use. Tourniquets were issued in Civil War surgical sets,<sup>5</sup> and the failure to apply one to a wounded Confederate Army general may have affected the outcome of that war and thus the course of U.S. history.<sup>6</sup> Paradoxically, it is perhaps from the same war that initial ambivalence to tourniquets arose: Prolonged time from tourniquet placement to definitive care often resulted in severe ischemic complications. This caused some surgeons of the day to argue that it was safer to allow continued bleeding than use a tourniquet to stop it.<sup>7</sup>

Tourniquets have long been standard issue in military medics' kits, yet there has been reluctance to use them in all but the direst of circumstances. In the 1960s, there were even efforts to have them removed from medical kits and deleted from the training curricula of military medics.<sup>2</sup>

Mabry describes a cycle wherein the tourniquet is initially welcomed by the military but soon falls out of favor due to perceived misuse, while many who might have been saved die of potentially controllable hemorrhage.<sup>7</sup> This cycle was repeated until analysis of mortality data from the Vietnam conflict led to renewed interest in the use of the tourniquet. This analysis suggested that a sizeable proportion of combat fatalities could have been averted by use of a tourniquet. In one report, it was estimated that 105 (38%) of

277 soldiers who died from extremity artery bleeding might have been saved by proper, timely tourniquet application.<sup>8</sup>

Recent experience has reinforced this trend toward more liberal military usage of tourniquets.<sup>9-11</sup> Modern combat results in high rates of extremity trauma. This fact, combined with the recognition that many who died of combat-inflicted wounds might have survived if a tourniquet had been used to arrest exsanguinations, forms the rationale for use of a tourniquet under modern combat conditions.<sup>12</sup> Sebesta, who has detailed his experiences as a surgeon in an Army Combat Support Hospital (CSH), states "tourniquets are an essential therapy based on recent experience in Iraq."<sup>9</sup>

This rationale is further reinforced by the circumstances under which much of the prehospital care is provided during a military conflict: Hostile action by adversaries, unfavorable environmental conditions, frequently prolonged transport to advanced care, austere logistics, and multiple casualties with limited triage and treatment manpower all support the expedient use of tourniquets on the battlefield. It is to be noted that these conditions are occasionally, if infrequently, experienced in civilian EMS settings as well.

The U.S. Army and Marine Corps both now issue tourniquets to individual soldiers and marines in their Individual First Aid Kits (IFAK) and train them in their proper use.<sup>13,14</sup> The Army prehospital trauma life support (PHTLS) mnemonic is now "MARCH" (Massive bleeding, Airway, Respirations, Circulation, and Head Injury) rather than "ABC" (Airway, Breathing, and Circulation), recognizing that massive hemorrhage on the battlefield is the primary treatable threat to survival and must be quickly arrested.<sup>9</sup>

Still, despite these changes, more than 50% of the deaths from isolated extremity hemorrhage were potentially preventable by correct tourniquet application, according to one report from the Iraq conflict.<sup>11</sup> It is not known how many patients with multisystem injuries had tourniquets placed nor if this intervention improved or worsened outcome.<sup>15</sup> Further reviews of battlefield tourniquet use will better document these results.

### Civilian EMS Usage

There has been a movement toward liberalization of tourniquet use in civilian EMS systems, but this remains controversial. Most systems still employ the tourniquet as a technique of last resort, using protocols that recommend direct pressure, pressure dressings, pressure points, elevation, and cold application as primary treatments for severe extremity hemorrhage. Of these, only direct pressure can be supported based on available evidence.<sup>4</sup> The most recent National Association for EMS Physicians (NAEMSP) consensus statement on wound care for delayed or prolonged transport

recommends the use of tourniquets *only* in cases of amputation.<sup>16</sup>

Despite extensive experience with tourniquets in the military medical services of the United States and other countries, recent civilian EMS teaching has not fully accepted this potentially life-saving instrument.<sup>17</sup> Indeed, many EMS systems do not allow their crews to carry tourniquets. Unfortunately, this can lead to circumstances in which a tourniquet is required and may have to be improvised. Improvised tourniquets are less likely to be effective<sup>10</sup> and may be more prone to neurovascular complications.

The positive experience with tourniquets on the battlefield holds promise for civilian EMS trauma care: Modern military protocols for tourniquet use could easily be incorporated into civilian EMS systems.

## Physiology, Complications, and Safe Use of Tourniquets

### Physiology

Arterial tourniquets work by compressing muscle and other tissues surrounding extremity arteries that, in turn, collapse the lumina of these arteries and thereby arrest flow distal to the tourniquet. The tension or force needed in order for a tourniquet to compress the artery is dependent on the size of the extremity as well as the width of the tourniquet. In general, larger circumference of an extremity correlates with higher required tension.<sup>18</sup> Wider tourniquets typically are more effective at stopping arterial flow at a given tension than narrow tourniquets.<sup>19</sup>

### Complications

Tourniquet use, which is well accepted as a technique for bloodless extremity surgery, has been associated with local and systemic complications (see Table 1).

TABLE 1. Potential Complications of Use of Tourniquets<sup>a</sup>

Local	Systemic
Postoperative swelling and stiffness	Increased central venous pressure
Delay in recovery of muscle power	Arterial hypertension
Compression neuropraxia	Cardiorespiratory decompensation
Wound hematoma	Cerebral infarction
Wound infection	Alterations in acid-base balance
Direct vascular injury	Rhabdomyolysis
Bone and soft-tissue necrosis	Deep venous thrombosis
Compartment syndrome	Tourniquet pain
	Systemic inflammatory response syndrome <sup>b</sup>
	Fibrinolysis <sup>c</sup>

Complications of operative tourniquets that have been reported in the surgical literature are presented here.

<sup>a</sup>Ref. 20.

<sup>b</sup>Ref. 31.

<sup>c</sup>Ref. 32.

Early surgical use of tourniquets led to the recognition that improper tourniquet design or prolonged tourniquet application (longer than 1.5 to two hours) could lead to muscle, nerve, and vascular injuries, resulting in a syndrome known as *tourniquet palsy* or *tourniquet paralysis*. Additionally, irreversible ischemic damage to limbs is known to occur in cases where a tourniquet had been left in place for longer than six hours; amputation of the limb above the level of the tourniquet was recommended in these circumstances and still remains a surgical dictum.<sup>2</sup>

*Tourniquet time* (i.e., the total time during which arterial flow beyond the instrument can be safely interrupted) is an issue of controversy. Evidence from animal studies shows that even minutes of tourniquet use will lead to changes in muscle and nerve physiology as well as systemic effects. These studies demonstrated that after one hour, there was no evidence of muscle damage, while two hours of ischemia led to elevated levels of both lactic acid and CPK, suggesting muscle damage was occurring.<sup>21</sup>

Most surgical guidelines recommend, and clinical studies support, no more than 60–90 minutes of operative tourniquet time in order to safely use this technique. Two hours of tourniquet time is a “useful guideline” for an upper limit.<sup>20</sup> Patients with advanced age, vascular diseases, and traumatic injuries are at higher risk for complications, including nerve and muscle injury. Nerve injuries have been reported after only 30 minutes of tourniquet time. Muscle, especially that directly under the tourniquet, has shown damage after one hour, though actual myonecrosis seems to occur only after three hours.<sup>22</sup> *Post-tourniquet syndrome* (comprising weakness, paresthesias, pallor, and stiffness) is common but seems to resolve after about three weeks.<sup>23</sup>

Recent military experience supports the safety of these short tourniquet times in prehospital patients.<sup>9–11</sup> Chambers et al. reported limb salvage in 11 of 14 (79%) of patients with arterial injuries despite total tourniquet times averaging two hours.<sup>24</sup>

All known complications of tourniquets seem to worsen with prolonged tourniquet time. Unfortunately, tradition has held that tourniquets, once placed, should be left on until removed by a physician. This tenet likely arose from the recognition that repetitively loosening and retightening a tourniquet exacerbates blood loss. While such “reperfusion intervals” are controversial and discussed in more detail below, our proposed protocol (and current military doctrine) allows for reevaluation of the need for, and possible removal of, a tourniquet by EMTs prior to reaching the hospital.

Tourniquet use may also result in venous complications, including worsened venous bleeding and venous thromboembolism (VTE). One major criticism of tourniquets is that, if not properly applied, tourniquets can actually increase bleeding by occluding venous return while not completely arresting arterial inflow.

Thrombosis could occur due to venous stasis during tourniquet use. Subsequent embolization of the clot(s) to the pulmonary circulation could then occur, either before or after tourniquet removal. The role of tourniquets in inducing venous thrombosis and pulmonary embolism is not clear. VTE has been reported to increase with tourniquet use in surgery; however, others have suggested that this complication is a result of surgery itself, not merely of tourniquet use.<sup>25–27</sup>

Elastic or compression dressings can be similarly criticized, as both may increase bleeding and promote venous stasis and VTE. A further drawback is that if placed too zealously, they can become an unrecognized arterial tourniquet.

Compartment syndrome has been a reported complication of tourniquet use. In most cases, this is felt to result from the injury necessitating tourniquet use, rather than the tourniquet itself, except in prolonged tourniquet-induced ischemia (more than three hours) or excessively high tourniquet pressures.<sup>23</sup>

Systemic acid-base changes may result from release of a tourniquet in place for an extended period of time. Limb ischemia results in lactic acidosis of tissues distal to the tourniquet. After release of the tourniquet, reperfusion of the extremity carries this acid and free radicals into the central circulation, a syndrome labeled *ischemia-perfusion injury*. Hyperkalemia and systemic acidosis may result in cardiac arrhythmias, among other problems. Clinical experience on the systemic metabolic effects of tourniquet release is inconsistent, and may vary with anesthetic technique. One study showed no such results after one to three hours of tourniquet time in a sample of elderly orthopedic surgery patients,<sup>28</sup> while a second showed that arterial pH, P<sub>a</sub>O<sub>2</sub>, P<sub>a</sub>CO<sub>2</sub>, lactate, and potassium changed significantly after tourniquet release.<sup>29</sup>

Hypertension and increased central venous pressure following operative tourniquet application are well-documented, but may be related to surgical practice: elevation and compression of the extremity to create a bloodless field results in autotransfusion of extremity blood into the central circulation.<sup>30</sup> Such an event is not likely to occur with field use of the tourniquet.

Other systemic changes, such as creation of a systemic inflammatory response and increased fibrinolytic activity, seem to be transient and are not known to be clinically significant.<sup>31,32</sup>

Pain from tourniquet use is a major concern. Some have stated that tourniquets can cause “excruciating pain”<sup>33</sup> despite proper application. In one report, however, awake, nonanesthetized volunteers who had tourniquets placed and inflated to 100 mmHg above their systolic blood pressure tolerated the instruments for 25 minutes on their forearms and 18 minutes on the upper arm.<sup>34</sup> It is not clear if a lower pressure, sufficient only to arrest bleeding, would be tolerated longer. Lower extremities, perhaps due to increased

circumference, have higher average times of pain tolerance, around 30 minutes.<sup>35</sup> It is clear, however, that most awake patients on whom a tourniquet is used will require medication for pain control.<sup>36</sup>

### Safe Prehospital Use

Safe prehospital tourniquet use depends on a number of factors. Underlying all safe prehospital tourniquet use are conservative and specific protocols defining indications, application and removal techniques, and application times. As always, regular training in protocols for prehospital providers is crucial. The outline of safe, effective protocols for prehospital tourniquet use can be extrapolated from both widespread surgical use and recent military experience.

The fundamental factors relating to safe tourniquet use are: tourniquet design, placement location, tourniquet tightness, and tourniquet time.

In terms of tourniquet design, it is well known that wider tourniquets with rounded, rather than sharp, edges are best in terms of limiting damage to underlying structures. The tourniquet should be made of a uniform, smooth material, as those with wire reinforcements are known to predispose to direct vascular injury due to unequal application of pressure under the wires.<sup>37</sup> As a pneumatic tourniquet, a blood pressure cuff is theoretically ideal, as it provides uniform pressure over a wide area. Its practical use, however, is somewhat limited by its size and weight, as well as its inability to maintain high pressures for prolonged periods. It is also difficult to apply securely to a short residual stump in the case of a traumatic amputation.

Most operative manuals recommend tourniquet placement on the thickest portion of the limb in order to maximize the tissue through which pressure is exerted and minimize the pressure required to stop arterial flow and thus the risk to underlying skin, muscle, nerves, and vessels. This may also limit the pain associated with tourniquet use, though some studies contradict this.<sup>33,38</sup>

EMS providers, however, have traditionally been trained to place the tourniquet just above the injury, while avoiding placement over a joint.<sup>39</sup> This more distal placement recommendation probably arises from concerns about the need for an amputation after definitive care is reached. The goal is to preserve as much limb length as possible. However, with proper tourniquet design and limited tourniquet time, a more proximal placement of the largest portion of the extremity is preferred because of speed of application, minimization of pressure injury to underlying tissues, and the possibility that multiple distal bleeding sites exist.

When applying a tourniquet, the lowest effective pressure should be used in order to minimize subsequent ischemic complications: A tourniquet must be

tightened only to the pressure required to arrest hemorrhage.

There is no rationale for using an occlusive tourniquet as a high-pressure dressing by placing it directly over a wound dressing, as it will not effectively stem arterial inflow to the wound in this location. Used nonocclusively, however, a tourniquet could effectively be used to augment a pressure dressing and hold it tightly in place, as suggested in a recent review of tourniquet use.<sup>40</sup>

Some have attempted to prolong tourniquet time by use of "reperfusion intervals." Although taught in some popular wilderness first-aid manuals,<sup>41,42</sup> these intervals are not practical in prehospital scenarios. These have been shown clinically to reduce complications only if perfusion is restored for 30 minutes or more.<sup>43</sup> Therefore, to be effective at reducing ischemic complications, they would likely also allow slow exsanguination. An interesting suggestion designed to lessen injury to tissues directly under the device is to use two adjacent tourniquets, alternately employing one then the other.<sup>44</sup>

All tourniquet usage must be well documented, then communicated on transfer of care. This minimizes the likelihood that a tourniquet will be overlooked by subsequent care providers and inadvertently left on for a prolonged period. Time of application must be recorded, either on the triage tag or physically written on the skin of the victim. The forehead is suggested as a prominent location. Triage cards should clearly annotate that a victim is wearing a tourniquet and the time of placement, as does the current DD Form 1380 *Field Medical Card*. One early advocate of the tourniquet recommended that casualties who are conscious be instructed to tell everyone with whom they come into contact that they have a tourniquet in place.<sup>45</sup> For the same reason, tourniquets should never be covered. There is evidence that cooling the extremity distal to the tourniquet may reduce complications.<sup>46</sup> A blanket placed over a tourniquet may be doubly dangerous, both warming the ischemic extremity and obscuring the tourniquet. The extremity is probably best left uncovered, except in temperatures where there is risk of direct cold injury. If available, a brightly colored marker may be placed at the tourniquet location as well.

The tourniquet should ideally be manufactured for its purpose. Improvised tourniquets will tend to apply pressure unevenly and often have sharp edges, increasing the risk of underlying tissue injury. Examples of suboptimal improvised tourniquets include belts and similar straps, which can entrap skin and directly cause injury. Cravats (i.e., triangular bandages) or elastic dressings (i.e., ACE<sup>®</sup> bandages) can bunch when twisted with a windlass.<sup>47</sup> Although these improvised options are frequently taught in first-aid manuals, they should be avoided unless no other options are available to arrest hemorrhage.

485 Finally, and most critically, tourniquet time must be  
 minimized. In most cases, this will mean transporting  
 the patient expeditiously to a higher level of care. In-  
 forming the transportation officer at a mass casualty  
 event of "tourniqueted" patients may allow these pa-  
 490 tients to have higher priority for transport to hospital.  
 In our protocol, no patient with a tourniquet will re-  
 ceive a triage code less acute than "yellow." In addition,  
 specific protocols can be utilized to remove tourniquets  
 that, upon reassessment at a later time, may no longer  
 be needed to control bleeding.

## 495 Alternatives to the Tourniquet

### Pressure Dressings

Pressure dressings are adequate to stop most cases of  
 hemorrhage, whether it occurs from the extremities or  
 other parts of the body. One commercially available  
 500 bandage already in use by the military for this purpose  
 is the "Israeli dressing." Other compression dressings  
 can be improvised with large amounts of gauze and an  
 elastic bandage that is wrapped around the wounded  
 505 limb, as described in a recent PHTLS manual.<sup>39</sup> These  
 bandages work on the principle of providing compres-  
 sion to reduce the flow of blood through damaged ves-  
 sels (primarily capillaries and veins) while providing  
 a "scaffold" on which blood can clot. As mentioned  
 510 previously, a carefully placed tourniquet could also  
 be utilized to tightly compress a bandage and act as  
 a pressure dressing, but only if it is not so tight as  
 to occlude arterial inflow or to increase distal venous  
 bleeding.

Experience with pressure dressings shows that they  
 515 work well in trained hands, stopping all "moderate  
 bleeding" and most "profuse bleeding" (81%), accord-  
 ing to one study. A tourniquet was required to stop one  
 case with "profuse bleeding."<sup>48</sup>

A major drawback of this type of dressing is that  
 520 they take time and often more than "one set of hands"  
 to apply properly. They must also be reassessed fre-  
 quently to ensure that bleeding has, in fact, been ar-  
 rested, which requires access to the site of the dressing  
 as well as a light source. It is foreseeable that circum-  
 525 stances may arise in which one or both of these are not  
 available, and time and personnel are needed to per-  
 form these assessments. In contrast, once placed and  
 tightened to arrest bleeding, modern tourniquets are  
 highly unlikely to lose effectiveness. After resuscitation,  
 530 however, they may require retightening if the victim's  
 blood pressure increases sufficiently to allow distal  
 flow.<sup>36</sup>

Self-application of an adequate pressure dressing can  
 be extremely difficult, if not impossible, and is cer-  
 535 tainly time-consuming. Modern tourniquets, including  
 the model preferred by the military, have been specifi-  
 cally designed for one-handed self-application. The goal

is rapid self-application in order to allow the victim to  
 continue the mission, if physically able to do so.

Finally, pressure dressings may be difficult or impos- 540  
 sible to secure to limbs that have sustained partial or  
 complete amputation.

In short, pressure dressings are excellent instruments  
 to control most causes of hemorrhage, but are not eas- 545  
 ily adaptable to circumstances in which there are lim-  
 itations on time and personnel, after amputation, or  
 when self-application is required. In these scenarios, a  
 tourniquet is much more easily and effectively used.

### Topical Hemostatic Agents

Topical hemostatic agents were developed in response 550  
 to the recognition that uncontrolled hemorrhage is  
 the major source of preventable mortality in com-  
 bat settings. Several have been deployed with U.S.  
 forces in recent combat actions around the world<sup>49</sup> and  
 their successes and shortfalls reported in the medical 555  
 literature.<sup>50-52</sup>

Although they may be useful adjuncts, these agents  
 do not have the same simplicity and effectiveness  
 of pressure dressings or tourniquets. Experience in  
 animal models has shown that many agents sim- 560  
 ply do not work quickly or well enough to stop  
 brisk bleeding.<sup>51</sup> HemCon is a gauze dressing impreg-  
 nated with chitosan (extracted from shrimp shells)  
 that assists in clotting. Although this dressing was  
 felt to work well for hemostasis on war-wounded 565  
 in Iraq, the majority of these were venous bleeds.<sup>52</sup>  
 Another agent, QuikClot, was shown to cause burns  
 and other soft-tissue complications in nearby tissue  
 when used in its initially marketed powder form.<sup>51</sup>  
 570 A newer formulation reportedly does not have this  
 complication.

Small, deeply penetrating wounds like those  
 produced by missiles are also problematic for topical  
 hemostatic agents, which must either be trimmed or  
 otherwise altered and placed in wounds in direct 575  
 contact with bleeding sites to ensure hemostasis. In the  
 case of the HemCon dressing, even large "cavitational"  
 wounds required that the dressing be placed under di-  
 rect vision, directly on the bleeding site, for the dressing  
 to be most effective.<sup>51</sup> Finally, it should be noted that, 580  
 with the exception of QuikClot, these are expensive,  
 perishable agents. These issues all limit these agents to  
 being useful adjuncts, rather than primary treatments,  
 for extremity hemorrhage in disaster and other EMS  
 585 settings.<sup>50</sup>

### Systemic Hemostatic Agents

Hemostatic agents that improve coagulation, espe-  
 cially in hypothermic, coagulopathic, and acidotic mul-  
 titrauma patients, have been in development and

590 preliminary use in the recent past. Factor VIIa con-  
centrate, for example, has been extensively used by  
surgical teams in Afghanistan and Iraq with favorable  
results.<sup>9,53</sup>

595 Little, if any, specific prehospital use of these agents  
has been reported.<sup>54</sup> In most cases, these agents have  
been used in patients with cavitory or visceral rather  
than external hemorrhage, and their applicability in  
extremity injuries is not known. In addition, current  
agents are prohibitively expensive for routine EMS and  
600 disaster use.

## Recommendations for Tourniquet Use in Civilian EMS Systems

### Tactical EMS and Police Officers

605 Tactical EMS (TEMS) providers, usually paramedics,  
support special weapons and tactics (SWAT) or special  
operations teams found in many civilian law en-  
forcement agencies, from the municipal to the federal  
level. Most of these personnel come from a background  
of civilian EMS agencies and are steeped in the tradi-  
610 tion of tourniquet avoidance. Current trends in law en-  
forcement, particularly as they impact on SWAT oper-  
ations, demand that we reevaluate the appropriateness  
of applying typical civilian EMS practices to law en-  
forcement and TEMS environments. Civilian firearms-  
615 related incidents are becoming more and more like mil-  
itary operations.

Military-style weapons with large, high-velocity bul-  
lets have proliferated in recent years as weapons of  
choice among gangs, narcotics traffickers, and terror-  
620 ist groups. No longer can law enforcement officers as-  
sume that they can easily "outgun" the suspects they  
are charged with apprehending.

Coincident with the change in the nature of these  
weapons is a change in the wounds and injuries result-  
625 ing from their use. Like today's military, police officers  
typically wear body armor as a means of mitigating  
some of the increased risk they now face from heavily  
armed suspects. While helping reduce risk of death or  
grave injury from torso wounds, these vests have not  
630 reduced the burden of extremity injury as a potential  
cause of death in urban conflicts.<sup>55</sup>

In 1997, a pair of heavily armed bank robbers wear-  
ing body armor held Los Angeles police at bay until  
being fatally wounded. One responding officer shot in  
635 the thigh reportedly lost an estimated 40% of his total  
volume of blood, lost consciousness, and nearly died.  
Had he had access to a tourniquet, he might have been  
able to stop the bleeding himself and perhaps even con-  
tinue to support the mission at hand.

640 The weapons used by police in responding to vio-  
lent criminals could also create circumstances in which  
the tourniquet may prove beneficial. A medical exam-  
iner stated one suspect in the event described above

died due to thigh wounds that caused exsanguination.  
Civilian EMS personnel were unable or unwilling to  
645 enter the scene.<sup>56</sup> The suspect's family sued the city of  
Los Angeles for this "notorious" inattention. Applica-  
tion of a tourniquet under these circumstances, perhaps  
even one given to the victim with instructions on how  
to place it, may have saved the life of this person.<sup>57</sup> 650

Beyond firearms, the use of explosives by terrorists  
and other elements is blurring the distinctions between  
military conflict and civilian crime. ATF and FBI statis-  
tics show that, even before the Oklahoma City bomb-  
655 ings and 9/11 made terrorism more of a concern in the  
United States, there were large numbers of bombings  
for criminal and "entertainment" purposes.<sup>58,59</sup>

The traditional civilian model of trauma care ex-  
emplified by the "golden hour" concept will blur  
along with these changes. Deaths from military-style  
660 weapons and explosives conform to a different distri-  
bution of time of death when compared with the more  
typical civilian trauma experience. Most deaths occur  
very early, prior to hospital care. Many of these are  
due to exsanguination<sup>60</sup> and extremity hemorrhage is  
665 the leading cause of death in potentially salvageable  
victims.<sup>12</sup>

These considerations all argue for training and equip-  
ping police officers for tourniquet use on themselves  
or others, much as the military has done for its front-  
670 line troops. While widespread tourniquet availability  
among "lay providers" may make some uneasy, we  
note that individual soldiers have been able to use  
the instrument successfully on themselves or wounded  
colleagues.<sup>15</sup> In addition, there is justification for allow-  
675 ing trained police officers to use tourniquets on civil-  
ian victims, as part of basic first aid. There is already  
widespread experience with police personnel deliver-  
ing first-responder care, including cardiac care via the  
use of AEDs. Police officers have also been trained to  
680 perform triage at mass casualty incidents.<sup>61</sup> Initial con-  
cerns about acceptance of these roles by police officers<sup>62</sup>  
have not been borne out.<sup>63</sup>

### Disaster Situations/Mass Casualty Triage

Isolated extremity injury causing exsanguination also  
685 occurs in civilian EMS practice. Preventable deaths  
due to failure of prehospital personnel and hospital  
providers to stop limb hemorrhage have been reported.  
In one study, 57% of those dying in metropolitan Hous-  
690 ton due to isolated penetrating extremity trauma had  
bleeding sites amenable to tourniquet therapy.<sup>64</sup>

The events of 9/11/2001 showed that global terror-  
ism can now be a local occurrence. With terrorism  
come weapons of mass effect. While biological (such  
as anthrax), chemical, and nuclear/radiologic weapon  
695 threats have received great focus, attacks using con-  
ventional weapons such as explosives and firearms still

prevail as "the most common type of terrorist attacks in modern history."<sup>65</sup>

700 Casualty data and mortality trends from civil unrest, especially bombings, have bolstered the arguments for use of tourniquets in prehospital care: penetrating extremity injuries occur in about half of the severely injured.<sup>66</sup> This increase in extremity trauma is not limited to adults: Pediatric victims of violence also have higher rates of penetrating extremity injury than do child victims of "non-terror-related injuries."<sup>67</sup>

705 Explosives, via primary, secondary and tertiary blast effects, induce amputations, partial amputations, and penetrating wounds of the extremities in bystanders. Recent experiences in Lebanon,<sup>68</sup> Palestine,<sup>69</sup> Israel,<sup>65</sup> Kosovo,<sup>70</sup> Bali,<sup>71</sup> Madrid,<sup>72</sup> London,<sup>73</sup> and other settings have shown that there can be large numbers of victims with complex injuries, including mangled extremities, amputations, partial amputations, and missile injuries, in addition to head, spine, and visceral injuries.

720 It has been noted that these types of events combine the severe mechanisms of injury typically associated with military combat with the short intervals from injury to rapid transport and definitive treatment, which are more characteristic of the civilian trauma experience.<sup>74</sup> Under these circumstances, many victims need only very simple interventions from EMS providers: The use of tourniquets for brief periods to limit blood loss and expedite transport would be a rational, and possibly life-saving, intervention.

730 Control of bleeding is beneficial to patient survival.<sup>75</sup> Indeed, even in those who survive despite massive hemorrhage, reducing blood loss and thereby preserving vital oxygen-carrying capacity will lessen complications, such as the adult respiratory distress syndrome and multisystem organ failure. A tourniquet that completely arrests hemorrhage before resuscitation will maximize preservation of red blood cells.

740 A triage officer or EMS crew responding to mass casualty event must be able to act quickly with simple interventions in order to maximize victim survival. Taking the time (and personnel) to apply pressure dressings could impair the smooth implementation of triage algorithms in mass casualty situations. As an example, the most recent PHTLS handbook recommends 10–15 minutes of direct digital pressure to stop bleeding.<sup>39</sup> Clearly, this will be impractical or impossible in many disaster situations. In addition to penetrating extremity injuries, head and torso injuries may also demand immediate stabilization. Multiple serious injuries make time-consuming hemorrhage control measures an unaffordable luxury.

750 In these circumstances, EMS personnel must have access to, and training with, arterial tourniquets. A protocol that allows tourniquets to be used as a first, rather than last, resort is imperative. Placing a tourniquet on a bleeding extremity, noting the time of placement, and

moving on to the next victim will allow providers to immediately stop bleeding that would otherwise contribute to hemorrhagic shock and may even cause fatality. Advanced Trauma Life Support (ATLS) training now acknowledges the need to stop "obvious external bleeding" during the primary survey.<sup>76</sup>

760 Once all victims have been triaged, or more help has arrived, prehospital personnel can return to those victims who have tourniquets in place. Wounds can then be reassessed and tourniquets possibly removed and replaced with pressure dressings. Although this approach seems to violate the traditional teaching that a tourniquet placed in the field must be left on until the victim reaches a hospital, there are circumstances in which this approach is reasonable, for example, if delayed or prolonged transport is anticipated. Algorithms have been developed for primary tourniquet placement and reevaluation with conversion to nontourniquet-based hemostasis<sup>10,77</sup> and successfully used in 76% of cases.<sup>10</sup>

775 Tourniquets are simple devices. Nonmedical safety personnel as well as lay people (such as "walking wounded") can be quickly trained to apply these devices safely and effectively. Victims themselves could even effectively use some types of tourniquets on their own wounded extremities.<sup>78</sup> Data from Canadian studies show that most commercial tourniquet models can be applied effectively in under 30–40 seconds.<sup>79</sup> Providers can be easily taught to quickly and effectively apply tourniquets. Life-support courses have been changing to simplify techniques for responders. Tourniquets should be taught as an adjunct to standard hemorrhage control techniques. Pressure points and elevation are commonly taught to lay persons. These techniques are arguably no simpler than tourniquet application and, unlike tourniquets, are of unproven benefit.<sup>4</sup>

### Routine EMS Usage

795 Tourniquets can also be useful in cases involving single patients, but only if EMS providers have access to appropriate protocols, training, and equipment. They must have familiarity with the indications and techniques for the use of tourniquets in order to avoid an inappropriate (and historically based) fear of these instruments. Tourniquets are naturally compatible with a "scoop and run" approach to trauma care in which simple, rapid, and potentially life-saving interventions are combined with expeditious transport to definitive care.

800 The use of a tourniquet to control extremity bleeding maximizes the ability of EMS providers to resuscitate a hypotensive patient by "stopping the leak." Ongoing extremity bleeding will hinder adequate resuscitation. Intravenous (IV) fluid infusions will simply dilute valuable oxygen-carrying hemoglobin and clotting factors.<sup>63,80</sup>

810 A "tourniquet first" approach to the single, multiply  
 injured patient allows the provider to immediately stop  
 obvious extremity bleeding, allowing attention to be  
 then turned to more time-consuming airway or breath-  
 ing priorities. After these are attended to, attention can  
 815 be directed to the bleeding extremity, with possible ex-  
 change of the tourniquet for a pressure dressing.

Transportation of prehospital patients is also facili-  
 tated by placement of tourniquets. Personnel do not  
 have to maintain digital pressure or frequently check  
 and reinforce dressings. This frees the provider to pay  
 820 more attention to maintenance of vital signs, to obtain  
 IV access, or to complete a secondary survey en route to  
 the hospital. In austere circumstances such as wilder-  
 ness settings, disasters, or hostile-fire situations, this  
 825 advantage is made even more significant as providers  
 are freed to perform other roles or to care for multiple  
 patients.

We also know that field-improvised tourniquets  
 may not be as safe as commercially manufactured  
 830 tourniquets (bandage/windlass combinations can  
 "bunch" into a constricting band<sup>47</sup>) and are frequently  
 ineffective.<sup>10</sup> It is better to have equipment, specifically  
 designed for the task of controlling catastrophic bleed-  
 ing in the hands of personnel well-trained in its use than  
 835 to ask them to fabricate crude devices under stressful  
 and possibly physically threatening circumstances.

## Protocols for Prehospital Tourniquet Use for Severe Extremity Hemorrhage

### Indications

840 Table 2 presents proposed indications for prehospital  
 tourniquet use, including routine EMS use in nondis-  
 aster settings. The goal of tourniquet use as presented  
 here is to allow prehospital personnel to safely, rapidly,  
 and effectively stop extremity hemorrhage, thus free-

TABLE 2. Indications for Tourniquet Use in Emergency Medical Services (EMS) and other Prehospital Settings

Amputation
Failure to stop bleeding with pressure dressing(s)
Injury does not allow control of bleeding with pressure dressing(s)
Significant <sup>a</sup> extremity hemorrhage in the face of any or all of:
Need for airway management
Need for breathing support
Circulatory shock
Need for other emergent interventions or assessment
Bleeding from multiple locations
Impaled foreign body with ongoing extremity bleeding
Under fire or other dangerous situation for responding caregivers
Total darkness or other adverse environmental factors
Mass casualty event <sup>b</sup>

Proposed indications for tourniquet use in EMS.

<sup>a</sup> "Significant" as defined by the EMS providers on scene.

<sup>b</sup> Any event where the number casualties and/or the severity of injuries exceed the ability of EMS personnel to provide optimal initial care all casualties.

ing the rescuer to triage and treat other patients or  
 rapidly address other emergent issues on a solitary pa- 845  
 tient. These principles are relevant to both mass casu-  
 alty situations and the care of a single patient. They  
 are intended to maximize the rescuer's efficiency and  
 enhance the safety of both patient and rescuer during 850  
 triage and treatment.

### Mass Casualty and Disasters

In the military environment, use of tourniquets during a  
 mass casualty event is well established and adaptation  
 of this use to civilian disaster care is natural.

Figure 1 shows an algorithm advocating that respon- 855  
 ders stop potentially massive bleeding first: this is easily  
 and safely accomplished with a tourniquet. The prompt  
 arrest of major extremity hemorrhage minimizes blood  
 loss while allowing the triage provider to move rapidly  
 to assess other patients. With multiple casualties, there 860  
 may not be the time or manpower to apply an adequate  
 pressure dressing.

After completing triage and other emergent proce-  
 dures such as airway stabilization, medics would then  
 be free to reassess the need for the tourniquet in pa- 865  
 tients on whom it was previously applied. Allowing  
 providers to reassess wounds for ongoing tourniquet  
 need under calmer circumstances maximizes the safety  
 and effectiveness of this tool. Tourniquet reassessment  
 and removal algorithms are presented below. 870

Finally, in order to minimize tourniquet times, we  
 recommend that no patient with a tourniquet in place  
 should have a triage code less acute than "yellow" and




FIGURE 1. Proposed algorithm for mass casualty tourniquet use. Triage teams should apply tourniquets to patients with bleeding extremity wound(s) and continue with START or similar triage protocols. They also mark or label the casualty to alert others to the tourniquet.




FIGURE 2. Proposed algorithm for routine EMS tourniquet use. Bleeding in patients with indications listed in Table 2 is arrested by a tourniquet placed primarily. If time, clinical situation, and personnel allow, a pressure dressing could be tried first.

that they have priority for transport. The presence of the tourniquet and the time that the tourniquet was applied should be prominently marked on each patient.

**Routine EMS Use and Law Enforcement Considerations**

Typical protocols for tourniquet application in EMS stringently limit the use of this device. In most simple single-patient extremity hemorrhage encounters, a tourniquet will not be required; standard pressure dressings will suffice. However, in cases of amputation, severe hemorrhage, multiple patients, or single patients with extremity hemorrhage combined with airway or breathing emergencies, a tourniquet must be available and providers must be well trained in its use. Figure 2 shows a proposed algorithm for safe and rational routine tourniquet use within a civilian EMS systems.

EMS providers are charged with providing sophisticated care in their daily jobs: airway interventions, vascular access, medication administration, defibrillation, and the like. Expecting them to be able to decide whether or not to apply a tourniquet is certainly within their scope of training and practice. Allowing them to reassess and remove tourniquets and monitor for further bleeding is also simply a matter of common sense, involving only simple protocols and training.

Control of bleeding is especially important in patients with multisystem trauma who need multiple interven-

tions and immediate transport. Rapid control of extremity bleeding with a tourniquet facilitates other interventions and allows rapid transport to definitive care while minimizing blood loss. In multiply injured patients, we suggest allowing providers to place a tourniquet *first* to stop blood loss immediately, then attend to airway, breathing, or other emergent priorities following the military's "MARCH" protocol. After these are addressed, the medic may then return to see if the tourniquet is still needed. In cases with short transport times, well within the known safety margins of tourniquet time, rapid transport without removal is indicated. When delayed or prolonged transport is anticipated, efforts to replace the tourniquet with a pressure dressing should be undertaken.

We also suggest that law enforcement officers, especially those in high-risk operational settings, be allowed to carry tourniquets and given training on how to use them on themselves, their teammates, and other victims, in an effort to stop severe extremity hemorrhage while awaiting EMS arrival.

**Reassessment and Removal of Tourniquets**

Previous discussions of prehospital tourniquets have typically recommended leaving tourniquets on until removal by a physician, regardless of the time involved, while some systems have advocated loosening tourniquets intermittently for brief reperfusion intervals in the event of prolonged transport. We reject both of these approaches. Based on available evidence, safe reassessment of tourniquet need and tourniquet removal in the field can be accomplished with simple, standardized protocols and training. Figures 3 and 4 show proposed algorithms for reassessing the need for and performing the removal of tourniquets, respectively.

**Future Directions**

**Improved Tourniquet Design**

Multiple design features to improve safety and effectiveness could be incorporated into tourniquets for use in prehospital and disaster situations. For example, curved tourniquets, which fit the natural conical taper of an extremity better than do straight rectangular tourniquets, provide hemostasis at lower pressures and seem to allow longer tourniquet times.<sup>20</sup>

Padding under a tourniquet with two-layer dressings, like stockinette or cast padding, reduces the skin damage these instruments can cause.<sup>81</sup> It may be possible to incorporate more padding into nonpneumatic tourniquet models and thus reduce the risk of skin injury. Other simple, potentially effective design changes would include widening the nylon strap on which most new models are based and coloring them brightly to make them conspicuous.

875

880

885

890

895

900

905

910

915

920

925

930

935

940

945

950




FIGURE 3. Steps to be followed to determine ongoing tourniquet need. The tourniquet should not be removed if the situation is unstable or there are not personnel or supplies available to place an adequate pressure dressing and monitor the site for rebleeding.

ously self-adjust tension as the victim’s blood pressure rises or falls, only applying the minimum pressure required to stop arterial flow. Other safety features, such as timers (allowing receiving facilities to know the tourniquet time) and alarms (to alert them to the tourniquet’s presence), might also be added.

**Tissue Protection**

Ischemia and reperfusion injury (IRI) is a known complication of prolonged tourniquet use and liberation of free radicals and other compounds after tourniquet release.

Recent work with n-acetylcysteine and preconditioning has not yet yielded success,<sup>82</sup> but as our understanding of this process improves, it is foreseeable that pre-hospital personnel will have access to compounds or techniques that may reduce IRI and further improve the outcome of tourniqueted limbs.

**Research and Registry**

Review of current and future uses of tourniquets is needed to continually improve our tourniquet protocols. Tumor boards, wound and trauma registries, and other prospective cohorts have enhanced medical knowledge about a number of conditions. Similar registries of tourniquet use should be encouraged and would be especially easy to initiate using data from recent military experience. This will help refine protocols, enhancing future safe tourniquet use. Controlled trials of prehospital tourniquet use are unlikely to be feasible given ethical and other clinical considerations. Appendix 1 presents a proposed data-collection form for use in a tourniquet registry.

**Pediatrics**

There is no reported prehospital experience with use of tourniquets for hemorrhage control in children. As previously noted, pediatric victims of violence often have the same injury patterns as adults; hence, they may also benefit from rapid extremity hemorrhage control with tourniquets. Further research to evaluate special considerations such as tourniquet size, childhood physiology, and other pediatric-specific issues must be performed in order to ensure maximum safety and benefit for all age groups.

**Training**

Continued safe and effective tourniquet use in the pre-hospital arena will be fostered by adequate training of personnel in use of this instrument. Appendix 2 presents a suggested outline of a tourniquet use training curriculum.

955

Finally, microelectronics could be incorporated to maximize both the effectiveness and the safety of tourniquets. Such “smart” tourniquets might detect flow through arteries beneath them and continu-




FIGURE 4. Algorithm for field tourniquet removal. A pressure dressing should be applied (\*with a topical hemostatic agent if allowed and available.) The tourniquet is loosened but left on and the wound monitored. If rebleeding occurs, tighten the tourniquet to arrest bleeding.

## CONCLUSION

1005

Traditionally, the risk-benefit calculus involved in the EMS use of tourniquets has been encapsulated in the phrase "lose a limb and save a life,"<sup>5</sup> because these devices have the potential to cause ischemic damage to a limb even when applied to stop life-threatening hemorrhage. Some authors, concerned with inappropriate use of tourniquets as well as reperfusion injury despite appropriate use, recommend prohibition. They state that there is no "exclusively clinical" reason to apply tourniquets, which should not be used except in the exigent circumstances of military or disaster situations.<sup>83</sup>

1010

1015

In contrast, Lee, et al. have suggested that there are "rare" circumstances when tourniquet use may be indicated.<sup>40</sup> However, recent military experience with widespread tourniquet use by individual soldiers, front-line medics, and combat hospital personnel, combined with almost universal acceptance of tourniquet use in bloodless extremity surgery, indicates that the maxim of "tourniquet as last resort" in civilian EMS care is clearly antiquated. Instead, there should be a prominent role for these potentially life-saving devices in civilian prehospital care. EMS providers must be trained and comfortable with tourniquet use when extremity bleeding is a threat and standard methods like direct pressure and elevation are ineffective or impractical.

1020

1025

1030

Penetrating extremity trauma is an increasingly common occurrence in our communities, whether associated with accidental injuries, firearm violence, or terrorist-related incidents. Experience and research indicate that life-threatening hemorrhage can be quickly and reliably arrested by the use of a simple tourniquet. This device allows limited numbers of providers to rapidly triage and provide hemostasis to multiple patients. Tourniquets can be self-applied by injured police, fire, or rescue personnel, allowing them to continue duty, if necessary, until safe evacuation and treatment are available.

1035

1040

Ischemic complications can be avoided by rational, protocol-driven use involving quick initial placement and rapid transport to definitive care, keeping tourniquet times to a minimum. When conditions allow, a tourniquet can be reassessed and replaced with a pressure dressing.

1045

## References

1050

1055

1. Coupland RM, Molde A, Navein J. Care in the field for victims of weapons of war: a report from the workshop organised by the ICRC on pre-hospital care for war and mine-injured. Geneva: International Committee of the Red Cross, 2001.
2. Klenerman L. The tourniquet in surgery. *J Bone Joint Surg Br.* 1962;44-B:937-943.
3. Welling DR, Burris DG, Hutton JE, Minken SL, Rich NM. A balanced approach to tourniquet use: lessons learned and relearned. *J Am Coll Surg.* 2006;203(1):106-15.

4. 2005 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care: Part 14: First Aid. *Circulation* 2005;112:IV196-IV203.
5. Kravetz RE. The tourniquet. *Am J Gastroenterol.* 2001;96(12):3425. Q4
6. Tarlowe MH, Swan KG. The death of Gen. Albert Sidney Johnson. *Bull Am Coll Surg.* 1998;83(5):15-23. 1065
7. Mabry RL. Tourniquet use on the battlefield. *Mil Med.* 2006;171(5):352-356.
8. Bellamy RF. The causes of death in conventional land warfare: implications for combat casualty care research. *Mil Med.* 1984;149(2):55-62. 1070
9. Sebesta J. Special lessons learned from Iraq. *Surg Clin North Am.* 2006;86(3):711-726.
10. Lakstein D, Blumenfeld A, Sokolov T, Lin G, Bssorai R, Lynn M, Abraham R. Tourniquets for hemorrhage control on the battlefield: a 4-year accumulated experience. *J Trauma* 2003;54(5-Suppl):S221-S225. 1075
11. Beekley A, Sebesta J, Blackburne L, Holcomb J. Pre-hospital Tourniquet Use in Operation Iraqi Freedom: Effect on Hemorrhage Control and Outcomes. Paper no. 27. Presented at Western Trauma Association Annual Meeting, March 2, 2006. 1080 Q5
12. Champion H, Bellamy R, Roberts P, Leppaniemi A. A profile of combat injury. *J Trauma* 2003;54(5-Suppl):S13-S19.
13. Statement by Lieutenant General Kevin C. Kiley, MD, Surgeon General of the United States Army. Medical Hearing, Committee on Appropriations, Subcommittee on Defense, United States Senate, First Session, 109th Congress, May 10, 2005. 1085
14. Suzuki S. 3/7 corpsmen go back to basics. *Marine Corps News*, p. 30 September 17, 2005. Accessed December 14, 2006, at: <http://www.usmc.mil/marinelink/mcn2000.nsf/main5/7780F210986B6FEB85257087004F21E8?opendocument>. 1090
15. Personal Communication with Major Alek Beekley, December 4, 2006.
16. NAEMSP Clinical Guidelines for Delayed or Prolonged Transport IV. Wounds [Prehosp Disaster Med. 1993;8(3):253-255] Reaffirmed 5/02. Accessed December 13, 2006. 1095 Q6
17. Varghese, M. Technologies, therapies, emotions and empiricism in pre-hospital care. In: Mohan D, Tiwari G, eds. *Injury Prevention and Control*. London: Taylor & Francis, 2000.
18. Shaw JA, Murray DG. The relationship between tourniquet pressure and underlying soft-tissue pressure in the thigh. *J Bone Joint Surg Am.* 1982;64(8):1148-1152. 1100
19. Crenshaw AG, Hargens AR, Gershuni DH, Rydevik B. Wide tourniquet cuffs more effective at lower inflation pressures. *Acta Orthop Scand.* 1988;59(4):447-451. 1105
20. Wakai A, Winter DC, Street JT, Redmond PH. Pneumatic tourniquets in extremity surgery. *J Am Acad Orthop Surg.* 2001;9(5):345-351.
21. Heppenstall RB, Balderston R, Goodwin C. Pathophysiological effects distal to a tourniquet in the dog. *J Trauma* 1979;19(4):234-238. 1110
22. Patterson S, Klenerman L. The effect of pneumatic tourniquets on the ultrastructure of skeletal muscle. *J Bone Joint Surg Br.* 1979;61-B(2):178-183.
23. Kam PC, Kavanagh R, Yoong FF. The arterial tourniquet: pathophysiological consequences and anaesthetic implications. *Anaesthesia* 2001;56(6):534-545. 1115
24. Chambers LW, Green DJ, Sample K, Gillingham BL, Rhee P, Brown C, Narine N, Uecker JM, Bohman HR. Tactical surgical intervention with temporary shunting of peripheral vascular trauma sustained during Operation Iraqi Freedom: one unit's experience. *J Trauma* 2006;61(4):824-830. 1120
25. Angus PD, Nakielny R, Goodrum DT. The pneumatic tourniquet and deep venous thrombosis. *J Bone Joint Surg Br.* 1983;65(3):336-339. 1125
26. Hirota K, Hashimoto H, Kabara S, Tsubo T, Sato Y, Ishihara H, Matsuki A. The relationship between pneumatic tourniquet time

- and the amount of pulmonary emboli in patients undergoing knee arthroscopic surgeries. *Anesth Analg.* 2001;93(3):776–780.
- 1130** 27. Jarrett PM, Ritchie IK, Albadran L, Glen SK, Bridges AB, Ely M. Do thigh tourniquets contribute to the formation of intra-operative venous emboli? *Acta Orthop Belg.* 2004;70(3):253–259.
28. Modig J, Kolstad K, Wigren A. Systemic reactions to tourniquet ischaemia. *Acta Anaesthesiol Scand.* 1978;22(6):609–614.
- 1135** 29. Townsend HS, Goodman SB, Schurman DJ, Hackel A, Brock-Utne JG. Tourniquet release: systemic and metabolic effects. *Acta Anaesthesiol Scand.* 1996;40(10):1234–1237.
30. Bradford EM. Haemodynamic changes associated with the application of lower limb tourniquets. *Anaesthesia* 1969;24(2):190–197.
- 1140** 31. Wakai A, Wang JH, Winter DC, Street JT, O'Sullivan RG, Redmond HP. Tourniquet-induced systemic inflammatory response in extremity surgery. *J Trauma* 2001;51(5):922–926.
32. Ellis MH, Fredman B, Zohar E, Ifrach N, Jedeikin R. The effect of tourniquet application, tranexamic acid, and desmopressin on the procoagulant and fibrinolytic systems during total knee replacement. *J Clin Anesth.* 2001;13(7):509–513.
- 1145** 33. Naimer SA, Chemla F. Elastic adhesive dressing treatment of bleeding wounds in trauma victims. *Am J Emerg Med.* 2000;18(7):816–819.
- 1150** 34. Maury AC, Roy WS. A prospective, randomized, controlled trial of forearm versus upper arm tourniquet tolerance. *J Hand Surg [Br].* 2002;27(4):359–360.
35. Hagenouw RR, Bridenbaugh PO, van Egmond J, Stuebing R. Tourniquet pain: a volunteer study. *Anesth Analg.* 1986;65(11):1175–1180.
- 1155** 36. Starnes BW, Beekley AC, Sebesta JA, Andersen CA, Rush RM Jr. Extremity vascular injuries on the battlefield: tips for surgeons deploying to war. *J Trauma* 2006;60(2):432–442.
37. Barnes JM, Trueta J. Arterial spasm: an experimental study. *Br J Surg.* 1942;30(117):74–79.
- 1160** 38. Finsen V, Kasseth AM. Tourniquets in forefoot surgery: less pain when placed at the ankle. *J Bone Joint Surg Br.* 1997;79(1):99–101.
39. Prehospital Trauma Life Support Committee of the National Association of Emergency Medical Technicians in cooperation with The Committee on Trauma of the American College of Surgeons. *PHTLS: Prehospital Trauma Life Support*, 6th ed. Edinburgh: Elsevier Mosby, 2006.
- 1165** 40. Lee C, Porter KM, Hodgetts TJ. Tourniquet use in the civilian prehospital setting. *Emerg Med J.* 2007;24(8):584–587.
- 1170** 41. Schimelpfenig T, Lindsey L. *Wilderness First Aid: National Outdoor Leadership School*, 3rd ed. Mechanicsburg, PA: Stackpole Books, 2000.
42. Weiss EA. *A Comprehensive Guide to Wilderness and Travel Medicine.* Oakland, CA: Adventure Medical Kits, 1997.
- 1175** 43. Horlocker TT, Hebl JR, Gali B, Jankowski CJ, Burkle CM, Berry DJ, Zepeda FA, Stevens SR, Schroeder DR. Anesthetic, patient, and surgical risk factors for neurologic complications after prolonged total tourniquet time during total knee arthroplasty. *Anesth Analg.* 2006;102(3):950–955.
- 1180** 44. Neimkin RJ, Smith RJ. Double tourniquet with linked mercury manometers for hand surgery. *J Hand Surg [Am].* 1983;8(6):938–941.
45. Tuttle AD. *Handbook for the Medical Soldier.* Baltimore, MD: William Wood and Company, 1927.
- 1185** 46. Swanson AB, Livengood LC, Sattel AB. Local hypothermia to prolong safe tourniquet time. *Clin Orthop Relat Res.* 1991;(264):200–208.
47. McLaren AC, Rorabeck CH. The pressure distribution under tourniquets. *J Bone Joint Surg Am.* 1985;67(3):433–438.
- 1190** 48. Naimer SA, Tanami M, Malichi A, Moryosef D. Control of traumatic wound bleeding by compression with a compact elastic adhesive dressing. *Mil Med.* 2006;171(7):644–647.
49. Shute N, Kelly K. Stopping the bleeding. Frontline medics in Iraq aim to treat the wounded better and faster. *US News World Rep.* 2003;134(11):30–31.
50. Pusateri AE, Holcomb JB, Kheirabadi BS, Alam HB, Wade CE, Ryan KL. Making sense of the preclinical literature on advanced hemostatic products. *J Trauma* 2006;60(3):674–682.
51. Wedmore I, McManus JG, Pusateri AE, Holcomb JB. A special report on the chitosan-based hemostatic dressing: experience in current combat operations. *J Trauma* 2006;60(3):655–658. **1200**
52. Wright JK, Kalns J, Wolf EA, Traweek F, Schwarz S, Loeffler CK, Snyder W, Yantis LD Jr, Eggers J. Thermal injury resulting from application of a granular mineral hemostatic agent. *J Trauma* 2004;57(2):224–230. **1205**
53. Martinowitz U, Kenet G, Segal E, Luboshitz J, Lubetsky A, Ingerslev J, Lynn M. Recombinant activated factor VII for adjunctive hemorrhage control in trauma. *J Trauma* 2001;51(3):431–438.
54. Holcomb J. The 2004 Fitts Lecture: Current Perspective on Combat Casualty Care. *J Trauma* 2005;59(4):990–1002. **1210**
55. Mabry RL, Holcomb JB, Baker AM, Cloonan CC, Uhorchak JM, Perkins DE, Canfield AJ, Hagmann JH. United States Army Rangers in Somalia: an analysis of combat casualties on an urban battlefield. *J Trauma* 2000;49(3):515–528. **1215**
56. Berry S, Glover, S Bank robber bled to death unnecessarily; shootout: accounts show mistakes by police, firefighters kept North Hollywood suspect from getting care. Officials give a different version. *Los Angeles Times* Los Angeles, California: April 21, 1998, p. 1. **1220**
57. Shuster, B Bank robber bled to death, autopsy finds; shootout: man was given no medical care and died of thigh wound in North Hollywood, report says. Whether accomplice was killed by police or self-inflicted gunshot is unclear. *Los Angeles Times* Los Angeles, California: April 11, 1997, p. 1. **1225**
58. Kapur GB, Hutson HR, Davis MA, Rice PL. The United States twenty-year experience with bombing incidents: implications for terrorism preparedness and medical response. *J Trauma* 2005;59(6):1436–1444.
59. Noji EK, Lee CY, Davis T, Peleg K. Investigation of Federal Bureau of Investigation bomb-related death and injury data in the United States between 1988 and 1997. *Mil Med.* 2005;170(7):595–598. **1230**
60. Gofrit ON, Leibovici D, Shapira SC, Shemer J, Stein M, Michaelson M. The trimodal death distribution of trauma victims: military experience from the Lebanon War. *Mil Med.* 1997;162(1):24–26. **1235**
61. Kilner T, Hall FJ. Triage decisions of United Kingdom police firearms officers using a multiple-casualty scenario paper exercise. *Prehosp Disaster Med.* 2005;20(1):40–46. **1240**
62. Groh WJ, Lowe MR, Overgaard AD, Neal JM, Fishburn WC, Zipes DP. Attitudes of law enforcement officers regarding automated external defibrillators. *Acad Emerg Med.* 2002;9(7):751–753.
63. Papson K, Mosesso VN, Jr. Ten years of police defibrillation: program characteristics and personnel attitudes. *Prehosp Emerg Care.* 2005;9(2):186–190. **1245**
64. Dorlac WC, DeBakey ME, Holcomb JB, Fagan SP, Kwong KL, Dorlac GR, Schreiber MA, Persse DE, Moore FA, Mattox KL. Mortality from isolated civilian penetrating extremity injury. *J Trauma* 2005;59(1):217–222. **1250**
65. Stein M, Hirshberg A. Medical consequences of terrorism. The conventional weapon threat. *Surg Clin North Am.* 1999;79(6):1537–1552.
66. Almogly G, Mintz Y, Zamir G, Bdolah-Abram T, Elazary R, Dotan L, Faruga M, Rivkind AI. Suicide bombing attacks: can external signs predict internal injuries? *Ann Surg.* 2006;243(4):541–546. **1255**
67. Aharonson-Daniel L, Waisman Y, Dannon YL, Peleg K, Members of the Israel Trauma Group. Epidemiology of terror-related versus non-terror-related traumatic injury in children. *Pediatrics* 2003;112(4):e280–e284. **1260**
68. Scope A, Farkash U, Lynn M, Abargel A, Eldad A. Mortality epidemiology in low-intensity warfare: Israel Defense Forces' experience. *Injury* 2001;32(1):1–3.

- 1265 69. Emile H, Hashmonai D. Victims of the Palestinian uprising (Intifada): a retrospective review of 220 cases. *J Emerg Med.* 1998;16(3):389-394.
70. Appenzeller GN. Injury patterns in peacekeeping missions: the Kosovo experience. *Mil Med.* 2004;168(3):187-191.
- 1270 71. Tran MD, Garner AA, Morrison I, Sharley PH, Griggs WM, Xavier C. The Bali bombing: civilian aeromedical evacuation. *Med J Aust.* 2003;179(7):353-356.
72. de Ceballos JP, Turegano-Fuentes F, Perez-Diaz D, Sanz-Sanchez M, Martin-Llorente C, Guerrero-Sanz JE. 11 March 2004: the terrorist bomb explosions in Madrid, Spain—an analysis of the logistics, injuries sustained, and clinical management of casualties treated at the closest hospital. *Crit Care.* 2005;9(1):104-111.
- 1275 73. Redhead J, Ward P, Batrick N. The London attacks—response: prehospital and hospital care. *N Engl J Med.* 2005;353(6):546-547.
- 1280 74. Shapira SC, Adatto-Levi R, Avitzour M, Rivkind AI, Gertsenshtein I, Mintz Y. Mortality in terrorist attacks: a unique modal of temporal death distribution. *World J Surg.* 2006;30(11):2071-2077.
- 1285 75. Kauvar DS, Lefering R, Wade CE. Impact of hemorrhage on trauma outcome: an overview of epidemiology, clinical presentations, and therapeutic considerations. *J Trauma* 2006;60(6 Suppl):S3-S11.
76. American College of Surgeons. *Advanced Trauma Life Support for Doctors—Student Course Manual*, 7th ed. Chicago, IL: American College of Surgeons, 2004. 1290
77. Walters TJ, Mabry RL. Issues related to the use of tourniquets on the battlefield. *Mil Med.* 2005;170(9):770-775.
78. Walters TJ, Wenke JC, Kauvar DS, McManus JG, Holcomb JB, Baer DG. Effectiveness of self-applied tourniquets in human volunteers. *Prehosp Emerg Care* 2005;9(4):416-422. 1295
79. King RB, Filips D, Blitz S, Logsetty S. Evaluation of possible tourniquet systems for use in the Canadian Forces. *J Trauma* 2006;60(5):1061-1071.
80. Fowler R, Pepe PE. Prehospital care of the patient with major trauma. *Emerg Med Clin North Am.* 2002;20(4):953-974. 1300
81. Olivecrona C, Tidermark J, Hamberg P, Ponzer S, Cederfjäll C. Skin protection underneath the pneumatic tourniquet during total knee arthroplasty: A randomized, controlled trial of 92 patients *Acta Ortho.* 2006;77(3):519-523.
82. Orban JC, Levraut J, Gindre S, Deroche D, Schlatterer B, Ichai C, Grimaud D. Effects of acetylcysteine and ischaemic preconditioning on muscular function and postoperative pain after orthopaedic surgery using a pneumatic tourniquet. *Eur J Anaesthesiol.* 2006;23(12):1025-1030. 1305
83. Husum H, Gilbert M, Wisborg T, Pillgram-Larsen J. Prehospital tourniquets: there should be no controversy. *J Trauma.* 2004;56(1):214-215. 1310

### APPENDIX 1. DATA COLLECTION FORM FOR TOURNIQUET REGISTRY

#### Tourniquet Registry Data Collection Form

1315 Date of Encounter: \_\_\_\_\_

Time of Dispatch: (use 24-hour clock for times) \_\_\_\_\_ Time of Arrival at Hospital: \_\_\_\_\_

Patient Identifier: (Run #, SSN, Med record, etc.) \_\_\_\_\_

1320 Patient Sex: M F (circle one) Patient Age \_\_\_\_\_ years old

Crew Members/Personnel: \_\_\_\_\_

1325 \_\_\_\_\_  
\_\_\_\_\_

Number of Victims on Scene: \_\_\_\_\_ Mechanism: Blunt Penetrating

1330 Site of Tourniquet Application: (circle; use additional sheets for > 1 tourniquet applied to single victim)

Arm Leg R L

1335 Tourniquet Applied by: Victim Bystander EMS Other: \_\_\_\_\_

Measures Used Prior to Tourniquet Use: (circle all that apply)

Direct Pressure Pressure Dressing Pressure Point(s) Hemostatic Agent

1340 Time of Tourniquet Application: \_\_\_\_\_ Time of Tourniquet Removal: \_\_\_\_\_

Tourniquet Removed by: (circle one) EMS (name \_\_\_\_\_) Hospital Personnel

1345 Total Tourniquet Time: (minutes) \_\_\_\_\_ Transport Time: (minutes) \_\_\_\_\_

Protocol Utilized for Tourniquet Placement: (circle one)

Mass Casualty/Disaster TEMS/Law Enforcement Routine EMS

1350 Removal In Field/En Route: (circle one)

Attempted/Successful Attempted/Failed Deferred

1355 Did Patient Require Pain Medications because of Tourniquet Pain? Yes No Unknown

Tourniquet-Related Complications (defined by higher level-of-care/Medical Control) (circle all that apply)

None Ischemic Damage VTE Compartment Syndrome Reperfusion Injury

1360 Other: (explain) \_\_\_\_\_

Type(s) of Bleeding Distal to Tourniquet: (as defined by higher level-of-care/Medical Control) (circle all that apply)

1365 Capillary Venous Arterial

**APPENDIX 2. SUGGESTED OUTLINE OF A TOURNIQUET-USE TRAINING CURRICULUM.**

**Tourniquet Training Curriculum Items:**

I. Background	1370
History of the Tourniquet Controversy	
II. Review of Hemorrhage Control	
Significance of Hemorrhage	
Hemorrhage Control Methods and Alternatives	1375
New trends: Military Use, Hemostatic Agents, etc.	
III. Protocols for Tourniquet Use	
Application	
Indications	1380
Mass Casualty/Disaster Situations	
TEMS/Law Enforcement	
Routine EMS	
Techniques	
Monitoring Effectiveness	1385
Removal	
Indications	
Techniques	
IV. Quality Improvement/Registry Instrument	1390
V. Practicum	
Scenarios for Practice Using Protocols	
Simulations to Practice Applying Tourniquet	
Self	1395
Partners	
Active Hemorrhage Simulators*	

\* from Mabry RL. Use of a hemorrhage simulator to train military medics. Mil Med. 2005 170(11):921-925.